

2007

Comment on "Intense Nonlinear Magnetic Dipole Radiation at Optical Frequencies: Molecular Scattering in a Dielectric Liquid"

Peter Palfy-Muhoray

Kent State University - Kent Campus, mpalfy@cpip.kent.edu

Follow this and additional works at: <https://digitalcommons.kent.edu/cpipubs>

 Part of the [Physics Commons](#)

Recommended Citation

Palfy-Muhoray, Peter (2007). Comment on "Intense Nonlinear Magnetic Dipole Radiation at Optical Frequencies: Molecular Scattering in a Dielectric Liquid". *Physical Review Letters* 99(18). doi: 10.1103/PhysRevLett.99.189401 Retrieved from <https://digitalcommons.kent.edu/cpipubs/237>

This Article is brought to you for free and open access by the Department of Chemical Physics at Digital Commons @ Kent State University Libraries. It has been accepted for inclusion in Chemical Physics Publications by an authorized administrator of Digital Commons @ Kent State University Libraries. For more information, please contact digitalcommons@kent.edu.

Comment on “Intense Nonlinear Magnetic Dipole Radiation at Optical Frequencies: Molecular Scattering in a Dielectric Liquid”

In the theoretical section of their Letter, the authors claim the result that “the magnetic current density is minus one half the total current density” [1]. In this Comment, I argue that their argument is flawed and their conclusion is invalid.

The authors consider an incident plane wave of angular frequency ω that propagates in the z -direction with its electric field linearly polarized along \hat{x} and its magnetic field oriented along \hat{y} . They then assert that, in conductors and insulators alike,

$$\nabla \times \mathbf{H} = \varepsilon_0 \dot{\mathbf{E}} + \mathbf{J}_c + \mathbf{J}_p + \mathbf{J}_M \quad (1)$$

where the \mathbf{J} 's are the conduction, polarization, and magnetic current densities. Integrating Eq. (1) over a surface perpendicular to the electric field gives

$$\int_S (\nabla \times \mathbf{H}) \cdot d\mathbf{s} = \int_S \varepsilon_0 \dot{\mathbf{E}} \cdot d\mathbf{s} + \int_S (\mathbf{J}_c + \mathbf{J}_p + \mathbf{J}_M) \cdot d\mathbf{s}. \quad (2)$$

The authors then invoke Faraday's law $\mathbf{H} = (\nabla \times \mathbf{E})/i\omega\mu_0$ and conclude that the integral on the left side of Eq. (2) is identical to the first integral on the right. The

theoretical result claimed in the Letter is directly predicated on this conclusion. This conclusion, however, is valid only because the authors use plane wave fields, which satisfy $\nabla \times \mathbf{H} = \varepsilon_0 \dot{\mathbf{E}}$ in the absence of currents. If, as the authors assert, there are bound and free electrons present, currents will be induced, and the fields will not be simply those of the incident plane wave. In this case, the two integrals will not be equal, and the conclusion will not be valid.

In summary, the authors' assumption of plane wave fields is inconsistent with the presence of currents. Their conclusion, that the magnetic current density is minus one half the total current density, is therefore invalid.

Peter Palfy-Muhoray
Liquid Crystal Institute
Kent State University
Kent, OH 44242, USA

Received 25 March 2007; published 2 November 2007

DOI: [10.1103/PhysRevLett.99.189401](https://doi.org/10.1103/PhysRevLett.99.189401)

PACS numbers: 42.65.-k, 32.10.Dk, 78.20.Ci, 78.20.Ls

- [1] Samuel L. Oliveira and Stephen C. Rand, Phys. Rev. Lett. **98**, 093901 (2007).