

7-2008

Comparative Chemical Versus Isotopic Dating of Monazites

R. Tracy

Virginia Polytechnic Institute and State University

C. Loehn

Virginia Polytechnic Institute and State University

Peter Dahl

Kent State University - Kent Campus, pdahl@kent.edu

M. Hamilton

University of Toronto

F. Mazdab

USGS

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.kent.edu/geolpubs>

 Part of the [Physical Sciences and Mathematics Commons](#)

Recommended Citation

Copyright 2008 Meteocritical Society.

This Conference Proceeding is brought to you for free and open access by the Department of Geology at Digital Commons @ Kent State University Libraries. It has been accepted for inclusion in Geology Publications by an authorized administrator of Digital Commons @ Kent State University Libraries. For more information, please contact digitalcommons@kent.edu.

Authors

R. Tracy, C. Loehn, Peter Dahl, M. Hamilton, F. Mazdab, J. Aleinkoff, and B. Miller

Comparative chemical versus isotopic dating of monazites

R. TRACY^{1*}, C. LOEHN¹, P. DAHL², M. HAMILTON³,
F. MAZDAB⁴, J. ALENIKOFF⁴, J. WOODEN⁴
AND B. MILLER⁵

¹Geosciences, Virginia Tech, Blacksburg, VA 24061
(*correspondence: rtracy@vt.edu)

²Geology, Kent State Univ., Kent OH 44242

³Univ. of Toronto, Toronto ON, Canada M5S 3B1

⁴USGS/Stanford SHRIMP Lab, Stanford CA 94305

⁵Texas A&M Univ., College Station TX 77843

Background

Monazites are widely used for U-Pb dating of igneous and metamorphic rocks and events. Monazite (mnz) U-Th-Pb dating that does not require isotopic data has recently been successfully utilized to characterize metamorphic events. In this “chemical” dating method, high-precision electron microprobe analyses with a significant spatial-resolution advantage over isotopic methods have been used to date *in situ* zones in partially recrystallized mnz grains that record timing of partial recrystallization events. This technique can produce a relatively complete picture of multiple tectono-thermal events in a metamorphic terrane or be used to decipher xenocryst-magmatic overgrowth relations in granitic plutons. The chemical method, however, remains controversial and in this presentation we report direct comparative results of *in situ* isotopic (SIMS, SHRIMP), bulk isotopic (ID-TIMS), and chemical (EMP) dating of the same mnz grains in rocks ranging from 2850 to about 300 Ma. We present analysis of uncertainties in chemical dating, including analytical errors and effects of common-lead correction.

Results

Case studies include (1) Archean and Paleoproterozoic mnz from the Tobacco Root Mountains, MT, (2) Mesoproterozoic mnz from a shear zone in NC, (3) Devonian mnz from granite, VT, and (4) Mississippian mnz from pegmatite, MA. In each case, dates are comparable within uncertainty, except where isotopic ages reflect mixing of two age zones.

Experimental calibration of zircon-quartz oxygen isotope fractionation

D. TRAIL^{1*}, I.N. BINDEMAN² AND E.B. WATSON¹

¹Department of Earth & Environmental Sciences, Rensselaer Polytechnic Institute, Troy, NY 12180, USA

(*correspondence: traild@rpi.edu)

²Department of Geological Sciences, University of Oregon, Eugene, OR, 97403, USA

Zircon is a common accessory phase occurring in many crustal rocks, and is often found with quartz. The $\delta^{18}\text{O}$ values of both minerals have been used to understand petrogenesis, and to constrain the provenance of grains no longer associated with their host rocks. In order to explore these minerals further, we report the first direct experimental characterization of oxygen isotope equilibrium fractionation.

Experiments (700-950°C; 10-20 kbar; 50-600 hrs.) were designed so that both phases would crystallize in chemical communication, but would be easily separable for isotope analysis. Zircon and quartz fractions were cleaned, and oxygen isotope analyses were performed on a Finnigan MAT 253 mass spectrometer by reacting zircon and quartz with BrF_3 using a 25W CO_2 laser line. Thus far, our data suggest an A-value which is in better agreement with empirical results [1] than with the indirect experimental calibration derived from zircon-water fractionation experiments [2].

A robust calibration would provide a new tool for mineral thermometry, and would have direct application to characterizing oxygen equilibrium/disequilibrium processes in petrological systems. In addition, because zircon is resistant to chemical alteration and physical breakdown during weathering, it is often used to constrain protolith origin(s). Our data provide an indirect means to determine $\delta^{18}\text{O}$ quartz values of the host rock – an important characterizing feature – provided zircon crystallization temperatures [3] and $\delta^{18}\text{O}$ zircon values are determined, and can be assumed not to have been compromised by post-crystallization diffusive exchange. The above methodology could also be applied to quartz inclusions in zircon which may be inaccessible to isotope analysis because of size, or altered by subsequent overprinting.

[1] Valley *et al.* (2003) *Geochim. Cosmochim. Acta.* **67**, 3257-3266. [2] Krylov *et al.* (2002) *European Journal of Mineralogy* **14**, 849-853. [3] Watson *et al.* (2006) *Contrib. Mineral. Petrol.* **151**, 413-433.